
HEALTH CONFIDENCE:

A SIMPLE, ESSENTIAL MEASURE FOR PATIENT ENGAGEMENT AND BETTER PRACTICE

Asking patients this one question can lead to better outcomes.

After being hospitalized for new onset congestive heart failure, Mrs. M, a 90-year-old widow, was discharged and sent home. Over the next six months, she had two additional hospital admissions and four emergency department visits without admission. She also had seven office visits and 21 telephone contacts with her primary care physician.

Eventually, her use of health care services dropped dramatically, and she returned to her usual activities. This turn of events came about because of a relatively simple intervention: Mrs. M's primary care physician asked her to assess her understanding of her condition and her confidence in managing it, and then worked with her to achieve gains.

"Naturally I was concerned about staying on my own when I came out of the hospital," she said. "At that time, my understanding of my heart failure was a '1' on a 10-point scale and my health confidence was a '4.' Now I am a '9' and an '8.'"

This case illustrates the downside of not paying attention to health confidence. Most of Mrs. M's post-discharge medical care was potentially avoidable. But when given the opportunity to gain insight and skills, she was able to better manage her condition and remain at home.

Consider another case:

Dr. A and colleagues observed very low health confidence and disease understanding among their many disadvantaged patients with hypertension. To address this, the clinic had its medical assistants use standard scripts to educate low-achieving patients during their visits. Within a year, more than 80 percent of these patients reported that they were very confident and understood their disease. As a result, blood pressure control improved significantly.


This second case demonstrates how proactive, "upstream" evaluation of health confidence can lead to interventions that improve patient outcomes.

Engagement and health confidence

A variety of terms – such as patient activation, patient self-efficacy, self-care, and self-management – have all been used to describe a patients' health-promoting actions. We prefer the term "engagement." The evidence-based, chronic care model illustrates the crucial connection between engagement and desirable patient outcomes.¹ For example, engaged patients have better

About the Authors

Dr. Wasson is an emeritus professor of community and family medicine and geriatrics at Dartmouth Medical School in Lebanon, N.H., and former director of the Center for Aging and the Dartmouth-Northern New England Primary Care Research Network. Dr. Coleman is director of the Care Transitions Program within the Division of Health Care Policy and Research at the University of Colorado at Denver, where he is also professor of medicine. Author disclosures: Dr. Wasson discloses that he is president of FNX Corp., which, under license with the trustees of Dartmouth College, makes the HowsYourHealth.org, HealthConfidence.org, and Medicare-HealthAssess.org tools described in this article available for free to office practices. Continuous users are asked to make a donation to support development and maintenance of the tools. Large health systems, employers, community sponsors, and office practices that exceed 3,000 uses a year are charged a nominal fee.



Research supporting the health confidence measure

health outcomes and better health care experiences, and likely use fewer health care services and cost less.²

Clearly, a patient's level of engagement is a good thing for a clinician and patient to know.

Health confidence is an effective proxy for engagement, and practices can easily measure it using a single question: "How confident are you that you can control and manage most of your health problems?" The possible responses are "very confident," "somewhat confident," "not very confident," and "I do not have any health problems." Alternatively, patients can rate their confidence on a scale from 0 (not very confident) to 10 (very confident). A score of

Several recent studies illustrate the clinical validity of the health confidence measure. In 69 primary care settings, patients' reported level of health confidence was the critical measure associated with a practice's performance outcomes.³ In the 23 clinical settings where health confidence levels ranked in the bottom one-third, 38 percent of patients exercised regularly, 73 percent engaged in healthy eating and risk avoidance, and 64 percent reported control of their blood sugar (if diabetic). In the 23 clinical settings where health confidence levels ranked in the top one-third, 75 percent exercised regularly,

ENGAGED PATIENTS HAVE BETTER HEALTH OUTCOMES AND BETTER HEALTH CARE EXPERIENCES, AND LIKELY USE FEWER HEALTH CARE SERVICES AND COST LESS.

7 or higher is the desired response. The question initiates self-reflection and meaningful communication between patients and health care providers. A helpful accompanying question is "How understandable and useful is the information your doctors or nurses have given you about your health problems or concerns?" (See "A tool to assess health confidence and begin action," page 10.)

Although we are using the term health confidence "measure" in this article, it should be thought of as more of an "action query" – that is, an assessment that results in immediate action and directly leads to improved patient engagement.

89 percent engaged in healthy eating and risk avoidance, and 78 percent reported control of their blood sugar.

Among 75,000 adults with chronic conditions, those who were very health confident reported many fewer adverse experiences than patients who were not as health confident.⁴ Patients who were not confident were much more likely to use the emergency department or be hospitalized and were more likely to claim that much of the hospital and emergency department use was unnecessary.

Patients' ratings of this single measure for health confidence are aligned with surveys based on multiple measures of their confidence to self-manage chronic conditions.⁴ ➤

In a cross-national study, 11 nations' rankings based on the single health confidence measure were strongly correlated with several measures of clinicians' engagement activities.⁴

Influences on health confidence

Practice patterns, comorbidities, and modes of administration can influence patients' reported levels of health confidence.

For both hospital and ambulatory care patients, health confidence is most positively associated with the quality of information

provided to the patient and access to care.^{3,5}

Low health confidence is strongly associated with low financial status and a high burden of illness.⁴ Persistent bothersome pain or emotional problems impede improvement in health confidence and undermine the effectiveness of care.⁶ Of Medicare beneficiaries studied who were health confident, more than 90 percent reported that they always take their medicines as prescribed.⁷ In contrast, only 75 percent who were not confident (and 60 percent who were not confident and also bothered by emotional problems) complied with medication therapy.

There is a crucial connection between patient engagement and desirable patient outcomes.

Health confidence is an effective proxy for engagement and can be measured by a single question.

Persistent pain or emotional problems can impede improvement in health confidence.

A TOOL TO ASSESS HEALTH CONFIDENCE AND BEGIN ACTION


Note: A pdf of this tool can be downloaded from the online version of this article: <http://www.aafp.org/fpm/2014/0900/p8.html>. Additionally, an electronic version and pdf versions in several languages are available at <http://www.healthconfidence.org>.

MY HEALTH CONFIDENCE

What number best describes your:

Health confidence

How confident are you that you can control and manage most of your health problems?




If your rating is less than "7," what would it take to increase your score?



Health information

How understandable and useful is the information your doctors or nurses have given you about your health problems or concerns?



If your rating is less than "7," what would it take to increase your score?



Family Practice Management[®]

Developed by John Wasson, MD, for HealthConfidence.org. Copyright © 2014 Trustees of Dartmouth College and FNX Corp. Used with permission. Physicians may duplicate for use in their own practices; all other rights reserved. <http://www.aafp.org/fpm/2014/0900/p8.html>.

Health confidence ratings are often positively biased if a patient is questioned directly by an authoritative figure, such as a health professional presenting the question with a judgmental tone.⁷ This is often referred to as the social desirability bias. To minimize it, the presenter should avoid a judgmental tone by emphasizing the value of an honest response.

Using the health confidence measure

Offices can implement the health confidence measure in several ways:

As a vital sign for patients who have chronic conditions, health risks, or difficulty following recommended care. Six of 10 adults who have chronic conditions are not health confident, and the majority of patients who have health risks (such as not exercising, being overweight, smoking, or drinking excessively) are not confident they will make

sary subsequent care.

For patients facing a new diagnosis, procedure, etc. Patients also experience significant transitions during clinic visits. New diagnoses, treatments, tests, and management decisions often disrupt a patient's health confidence.

If a patient's health confidence is high (a score of 7 to 10) and the patient has a good understanding of his or her health problems, the office can proceed with usual care.

If a patient's health confidence is low (a score of less than 7), the clinician or a staff member should follow three basic steps of motivational interviewing:⁹

1. Find a way to support or reinforce any progress the patient has made. For example, "I see you marked a '4' on health confidence. What made you mark a '4' instead of a '0'?" This allows the patient to reflect on some personal strengths.

Offices can use the health confidence measure as a vital sign for patients with chronic conditions or health risks.

For patients transitioning from another care setting, asking about health confidence can identify those who feel unprepared to manage their health problems.

If a patient's health confidence is less than 7, motivational interviewing techniques can help identify ways to improve the patient's score.

NEW DIAGNOSES, TREATMENTS, TESTS, AND MANAGEMENT DECISIONS OFTEN DISRUPT A PATIENT'S HEALTH CONFIDENCE.

progress in the two months after they choose to reduce one of these risks.⁸ To identify these patients and help them get the resources they need, medical assistants can routinely ask patients about their health confidence as part of their collection of vital signs.

As part of a routine check up. For example, the Medicare annual wellness visit offers a scheduled opportunity to integrate health confidence into a patient's health check up. Both paper-based and electronic assessment methods are available. (See the Medicare annual wellness visit tools listed under "Additional resources" on page 12.)

For patients transitioning from another care setting. Depending on their burden of illness and the quality of their hospital, 12 percent to 67 percent of patients feel prepared to manage their health problems at discharge.⁵ Exposing inpatients to the health confidence tool could help them express their limits and concerns as they transition from the hospital and could obviate much unneces-

2. Ask what it would take to reach a higher level of health confidence. For example, "What would you need to move from a '4' to a '7'?" This allows the patient to identify behavioral goals that are acceptable, meaningful, and feasible rather than having the clinician suggest a goal that does not fit with the patient's values.

3. Ask which goals might be most important and how the doctor or office staff might be able to help. This allows the patient to prioritize options, and allows the practice to be clear about the support it will be able to provide. Often, patients need a plan that involves Internet or community resources that teach them about problem solving, weight loss, blood sugar control, etc. (For more information on using community resources, see "Caring for Seniors: How Community-Based Organizations Can Help," page 13.) The support plan can be revisited, reinforced, and modified at subsequent visits.

When a patient's progress is slow, consider

whether pain or emotional problems (anxiety, depression, etc.) are present. An additional intervention such as telephone outreach is often needed to help these patients achieve behavior change. The patient's response to the question "What will it take to move higher on the confidence scale?" should guide the choice of resources. The patient may need to be referred to a health coach, case manager, or a specialist in behavioral therapy. Peers who have completed a chronic disease self-management program or those who are members of community support groups may be valuable resources as well.

Reassessment and reinforcement is usually needed to improve health confidence. When health confidence becomes part of everyday clinical service, the practice will find that it is a powerful stimulus for behaviorally sophisticated actions and a useful organizing principle for more efficient and effective services.

The next "blockbuster drug"

Patient engagement has been rightfully likened to a blockbuster drug for successful management of chronic conditions and the reduction of health risks.¹⁰ However, its appropriate prescription has been limited by its vague definition. Health confidence provides a free, easy to use, actionable measure to encourage patient engagement and better practice. **FPM**

1. Bodenheimer T, Lorig K, Holman H, Grumbach K. Patient self-management of chronic disease in primary care. *JAMA*. 2002;288(19):2469-2475.
2. Hibbard JH, Greene J. What the evidence shows about patient activation: better health outcomes and care experiences; fewer data on costs. *Health Aff*. 2013;32(2):207-214.
3. Wasson JH, Benjamin R, Johnson D, Moore LG, Mackenzie T. Patients use the Internet to enter the medical home. *J Ambul Care Manage*. 2011;34(1):38-46.
4. Wasson JH. A patient-reported spectrum of adverse health care experiences: harms, unnecessary care, medication illness, and low health confidence. *J Ambul Care Manage*. 2013;36(3):245-250.
5. Lepore M, Wild D, Gil H, Lattimer C, Harrison J, Woddor N, Wasson JH. Two useful tools to improve patient engagement and transition from the hospital. *J Ambul Care Manage*. 2013;36(4):338-344.
6. Wasson JH, Johnson DJ, Mackenzie T. The impact of primary care patients' pain and emotional problems on their confidence with self-management. *J Ambul Care Manage*. 2008;31(2):120-127.
7. Wasson JH. Unpublished national data obtained from 5,000 Medicare beneficiaries responding to MedicareHealthAssess.org, 2012-2014.
8. Wasson JH. Regular exercise is strongly associated with anticipated success for reducing health risks. *J Ambul Care Manage*. 2014;37(3):273-276.
9. Stange KC, Woolf SH, Gjeltema K. One minute for prevention: the power of leveraging to fulfill the promise of health behavior counseling. *Am J Prev Med*. 2002;22(4):320-332.
10. Dentzer S. Rx for the "blockbuster drug" of patient engagement. *Health Aff*. 2013;32(2):202.

Send comments to fpmedit@aafp.org, or add your comments to the article at <http://www.aafp.org/fpm/2014/0900/p8.html>.

ADDITIONAL RESOURCES

The following resources can assist practices with assessing patients and their health confidence:

HealthConfidence.org

<http://www.healthconfidence.org>

This site provides a complete health assessment with tailored information and a personal health plan for people age eight to 99. It also offers figures and forms related to this article, such as the "Problem Solving" tool (<http://www.healthconfidence.org/pblmslv/>), which can help patients prioritize their problems and develop practical solutions.

MedicareHealthAssess.org

<http://www.medicarehealthassess.org>

This site incorporates health confidence into a paper or electronic checkup for Medicare annual wellness visits. There is a "short form" version and a "long form" version, which automatically adds a personal health plan that the user can share and update at any time.

VeryIll.org

<http://www.veryill.org>

This site offers a tool for clarifying and supporting the needs of patients who are frail or need palliation or hospice care.

Health Assessments in Primary Care: A How-to Guide for Clinicians and Staff

<http://1.usa.gov/1p5wmGU>

This is a "how to" manual for health assessment in primary care practice created by the Agency for Healthcare Research and Quality. It offers additional links to information and communication technologies.

Patient Activation Measure

<http://bit.ly/OGd1ha>

This site offers a multi-item measure for patient activation that includes many items for patient health confidence.

Further reading:

Stewart EE, Fox C. Encouraging patients to change unhealthy behaviors with motivational interviewing. *Fam Pract Manag*. 2011;18(3):21-25. <http://www.aafp.org/fpm/2011/0500/p21.html>.

Hibbard J, Lorig K. The dos and don'ts of patient engagement in busy office practices. *J Ambul Care Manage*. 2012;35(2):129-132.

Simmons L, Baker NJ, Schaefer J, Miller D, Anders S. Activation of patients for successful self-management. *J Ambul Care Manage*. 2009;32(1):16-23.